






Pre-Op Information:
Shoulder Replacement Surgery at GBMC

GBMC Joint & Spine Center



Sign-up for GBMC MyChart

- My Chart allows you to verify your health insurance information, mailing address, phone number, e-mail, pharmacy, etc.
- MyChart will send reminders for your pre-op appointments and appointments while you are in the hospital.
- MyChart gives you access to your test results, allows you to communicate with your providers, pay your bill, and helps you to coordinate your healthcare needs.
- On the day of admission: MyChart will convert to MyChart Bedside, it is your portal for engaging with your care while admitted to the hospital. Empower yourself and your family with access to your care team, clinical data, and health education.







Shoulder Replacement Preop Checklist

		443-849-6261	-
		Surgeon:	
Place an X when task is completed Date of your surgery:			
[)	Activate your GBMC MyChart	Online link in Welcome E-mail
()	Review the Shoulder Replacement Presentation	http://www.gbmc.org/jointandspine
ſ]	Complete: Discharge Questionnaire	YES NO (MyChart or Packet)
_	ĺ	I-PSS (Male patient)	YES NO (MyChart or Packet)
[)	Do you know how to use your Hibiclens CHG wash?	Total of 4-CHG showers at home
•		The Antimicrobial Skin Cleanser, Disposable cloths, Instructions	Begin to use 3-days BEFORE surgery
		Pre-op MSSA/MRSA test at	Walk-in, no appointments needed.
٢)	GBMC's Diagnostic Testing Center	Monday-Friday
		Must be completed within 10 to 30 days	8 AM - 4 PM
		Must be completed within 10 to 30 days before surgery date.	8 AM - 4 PM no holidays
٢	1	61 1631	no holidays Call your surgeons office for
()	before surgery date.	no holidays Call your surgeons office for prescriptions issues, pharmacy
()	before surgery date. Discussion with ECIP Ortho Care Coordinator	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from
())	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!!
())	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from
(°)	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900 Date:Time: If you are missing any of these items, need assistance, or surgery is canceled: GBMC Joint & Spine Center	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from your pharmacy before surgery!! Please call 443-849-6261
)	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900 Date:Time: If you are missing any of these items, need assistance, or surgery is canceled:	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from your pharmacy before surgery!! Please call
)	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900 Date:Time: If you are missing any of these items, need assistance, or surgery is canceled: GBMC Joint & Spine Center	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from your pharmacy before surgery!! Please call 443-849-6261
)	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900 Date:Time: If you are missing any of these items, need assistance, or surgery is canceled: GBMC Joint & Spine Center 6701 N. Charles Street, Unit 58: Suite 5835, Towson MD 21204	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from your pharmacy before surgery!! Please call 443-849-6261 April or Casey
)	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900 Date:Time: If you are missing any of these items, need assistance, or surgery is canceled: GBMC Joint & Spine Center 6701 N. Charles Street, Unit 58: Suite 5835, Towson MD 21204 Shoulder Sling call Synergy Orthopedics 443-681-8971	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from your pharmacy before surgery!! Please call 443-849-6261 April or Casey Call today schedule a fitting:
	Notes	before surgery date. Discussion with ECIP Ortho Care Coordinator Dr. Lanzo'a patients: Victoria Karnas 443-849-3828 Dr. Buchalter's patients: 410-377-8900 Date:Time: If you are missing any of these items, need assistance, or surgery is canceled: GBMC Joint & Spine Center 6701 N. Charles Street, Unit 58: Suite 5835, Towson MD 21204 Shoulder Sling call Synergy Orthopedics 443-681-8971	no holidays Call your surgeons office for prescriptions issues, pharmacy location updates!! Pick up your medication(s) from your pharmacy before surgery!! Please call 443-849-6261 April or Casey Call today schedule a fitting:

Please review the
Joint & Spine Center
Pre-op Packet
for important information,
including this pre-op
checklist seen here.









Joint Replacement Surgery

Presented by: April Asuncion Higgins, BSN, RN, CMSRN

Time Sensitive Tasks to Complete Before Your Surgery

- ➤ Nasal swab for MRSA/MSSA
 - ➢ Golden window for completion is within 10-30 days prior to surgery. Must be completed no later than 10 days prior to your surgery date.
 - > Non-Kaiser Permanente patients: GBMC Ortho, GBMC Neurosurgery, Ortho Maryland
 - ➤ Testing at GBMC Diagnostic Testing Center
 - ➤ Monday-Friday 8:00 am 4:00 pm, no holidays
 - ➤ Results valid for 60-days
- ➤ Preop CHG Wash Kit with directions
 - ➤ Begin 3 nights before surgery
- ➤ Pre-op Questionnaires
- Fill your prescriptions before your surgery date
- > Pre-op Clearance from your Primary Care team(s), Diagnostic Tests, and blood work







Near EAST PAVILION-across the hallway from the Einstein Bagel Coffee shop









MON-Kaiser Permanente patients only Diagnostic Center and Laboratory Medicine at GBMC

GBMC Main Hospital 6701 N. Charles St./ Suite 3100/ Towson, MD 21204 MSSA/MRSA preop testing: Monday-Friday 8am-4pm, no holidays











Reasons for Joint Replacement Surgery

- ➤ Improve quality of living
- ➤ Tired of living in pain
- ➤ Not being able to do things you like to do

Shoulder Arthroplasty











https://orthoinfo.aaos.org/e n/treatment/reverse-totalshoulder-replacement/





Preoperative Testing & Plan of Care for Discharge

- Ortho Care Coordinator
 - ➤ Assists with scheduling Pre-op PT Home Visit for qualifying patients
 - ➤ Discharge planning begins
 - Home with outpatient physical therapy
 - ➤ Home with home physical therapy
 - > Subacute rehab
- ➤ Designating family/friends support person(s), and preparing your home environment

- ➤ MSSA/MRSA nasal swab to be completed 10 days to 30 days prior to your surgery date:
 - ➤ GBMC Diagnostic Testing Center:
 - ➤ Monday-Friday 8:00 am- 4:00 pm, NO HOLIDAYS
 - ➤ Kaiser Permanente patients are to have their pre-op MRSA collected at the Kaiser PEEC Center
- Need Surgical Clearance from
 - ➤ Primary Care Physician
 - ➤ Specialist, if applicable
- Your primary care physician office:
 - EKG, blood work, urinalysis, x-rays, etc.

Results and Surgical Clearance statements from you doctor(s) will need to be available to GBMC *no later than 3-days* before surgery.



F

How to Prevent Infection and Complications

- Control your blood sugar before, during and after your surgery
- Exercising and healthy weight loss
- ➤ You must STOP smoking (including cigars, e-cigarettes and vaping) 2 weeks before surgery and at least 2 weeks after surgery
- Good nutrition with adequate calories and protein
- Making sure skin is dry in between skin folds
- Frequent UTI's, notify your primary team & surgeon
- ➤ Advised no dental work within the 2weeks prior to surgery and/or the first 3-months after surgery.
- No artificial nails, acrylics, gel/dip/shellac polish on fingernails. Only clear, clean nails.

- Keep pets and animals away from the incision site after surgery.
- Applying fresh linens to bed before surgery. No pets are to sleep in bed or on the sofa with you until cleared by your surgeon.

PREVENTION









healthy eating

exectse

no smoking

glucose control







4 Days of Pre-surgery CHG Showers



https://youtu.be/eF3tae-c6d8

Hibiclens Shower Video


Please note **GBMC J&S Center**requires patients to complete 4
showers at home before surgery

- ➤ Pre-Surgical Scrub: 4% Chlorhexidine gluconate (CHG).
- > Four (4) Total CHG Showers at home.
- ➤ Begin three (3) nights before surgery, last application is on the morning of surgery before coming to the hospital.
- ➤ Apply the CHG foam onto your 6 or 7 disposable cloths, lather the solution onto skin.
- ➤ Avoid contact with your eyes, ears, mouth, and genitals.
- ➤ Do NOT shave or use hair removal lotions, deodorant, perfume, lotion, creams, or oils on your body.





Preventing Surgical Site Infection: Pre-Op Skin Preparation



- Hibiclens Shower #1 begin three (3) nights before your surgery
- ➤ Hibiclens Shower #2 is two (2) nights before surgery.
- Hibiclens Shower #3 is the night before surgery.
- Hibiclens Shower #4 is the morning of surgery before coming to the hospital.

Check off each yellow circle after your CHG shower



Hibiclens Shower Video
Please note GBMC J&S Center
requires patients to complete 4
showers at home before surgery







How It Works

Increased Protection Against Risk of Infections

The nose is a reservoir for bacteria that escape typical hygiene protocols such as hand washing. Nozin® Nasal Sanitizer® antiseptic equips health professionals, patients and individuals with a safe and effective nasal shield that helps decrease bacteria on nasal vestibule skin. Importantly, Nozin Nasal Sanitizer antiseptic is olinically shown to reduce nasal bacteria carriage without antibiotics.







Nozin® Nasal Sanitizer® is specially formulated to effectively address nasal carriage of bacteria:

- Effective kills 99.99% of germs and is clinically proven to decrease nasal bacteria carriage
- Safe, Fast-Acting the active ingredient is ethyl alcohol, a well-established, trusted, broad spectrum antiseptic
- 12-Hour Duration soothing solution is well tolerated, safe for regular use and Nasal Sanitizer® effect lasts 12 hours

- No Antibiotics alcohol-based antiseptic
- Pleasant experience a soothing, moisturizing feeling and a soft smell of citrus
- Easy To Use with the multiple use bottle, application of Nozin is simple and convenient

Do you have an *orange fruit oil allergy*?

Think hand sanitizer for your nose.

"To every patient, every time, we will provide the care that we would want for our own loved ones."





Coach/Support Person Enlist a family member or friend for help

- > Exercises
- ➤ Transfers to the toilet, tub, etc.
- ➤ Moving around with your walker or cane
- ➤ Bathing/dressing (using adaptive equipment if needed)
- ➤ Transportation to appointments
- ➤ Clean and prepare your home
- ➤ Food, shopping, preparing meals
- ➤ Cleaning laundry
- ➤ Caring for pets, children, grandchildren











Home Set-up

Before coming to the hospital

- Prepare meals ahead of time or arrange to have family/friends there to help
- Make frequently used items in the kitchen accessible for use
- > Clean and clear home of clutter
 - Remove rugs out of bathroom and kitchen
- ➤ Make sure all equipment that you may already have is easily accessible, clean and in safe working order
 - Borrow from friends or family or community loan closets









Can continue until day of surgery

- ➤It is OK to take:
 - ➤ Tylenol (acetaminophen)
 - ➤ Oxycodone, Tramadol, Dilaudid, Neurontin, Flexeril and Skelaxin
 - ➤ Celebrex is the only NSAID that is OK
 - ➤ Beta blocker and/or Calcium Chanel blocker
 - > Non-MAIO Anti-depressants, Anti-anxiety medications, Psychiatric medications
 - >Anti-Seizure medications
 - > Parkinson's medications
 - ➤ Asthma Inhalers/medications
 - ➤ Thyroid medications
 - >Steroids: oral or inhaled
 - ➤ Cholesterol Medication-Statins: Zocor, Lipitor, Pravastatin
 - **>**Birth Control Pills
 - **≻**Eye Drops
- >Get specific instructions from you provider for medications listed below:
 - ➤ Eliquis, Coumadin, Pradaxa, Plavix, Xarelto, Ticlid, Lovenox injections, Aspirin, or other blood thinners





10 Days before surgery

- ➤ Stop taking aspirin or salicylates, including Excedrin, Alka-Seltzer, Pepto-Bismol, unless otherwise directed by your provider
 - ➤ It is important to stop as they cause excessive bleeding during your surgery and recovery period.
- ➤ Stop all Herbal Products/Alternative Medications:
 - Examples: Vitamin E, glucosamine, fish oils, ginkgo biloba, ginseng, dong quai, garlic, turmeric, CoQ10, kava, ma huang, St. John's wort and non-vitamin supplements





7 Days before surgery

- > Stop NSAIDs products
 - ➤ Non-Steroidal Anti-inflammatory Drugs
 - Examples: Aleve, Ibuprofen (Motrin, Advil), Naproxen, Treximet, Vimovo, Duexis, Nabumetone, Ketolorac, Mobic, Robaxin, Meloxicam, Diclofenac
- > Stop all diet and weight loss medications
 - ➤ Examples: Phentermine, Glucomannan, CLA (conjugated linoleic acid), Orlistat (Alli), Garcinia cambogia, Hydroxycut, Raspberry ketones, Meratrim, green coffee bean extract, green tea extract, Forskolin, Synephrine, etc





3-days (72 hours) before surgery

- ➤ Stop Viagra, Levitra & Cialis
- >Stop topical medications



48 Hours before surgery

- ➤ **Stop** Monamine Oxidase Inhibitors (MAIOs)
 - Examples: Nardil (Phenelzine), Emsam (Selegline), Marplan (Isocarboxazid), Parnate (Tranylcypromine)





Do not take the day of your surgery

- **>Stop** ACE Inhibitors:
 - ➤ Examples: Lisinopril, Vasotec
- ➤ **Stop** Angiotensin Receptor Blockers:
 - Examples: Cozaar, Iosartan, Benicar, Candesartan, Atacand, Valsartan



F

Current Diabetic Medications

4 days before surgery

➤ Stop SGLT-2 Inhibitor medications: Steglatro

3 days before surgery

➤ Stop SGLT-2 Inhibitor medications: Jardiance, Invokana, Farxiga, and

24 Hours before surgery

- ➤ Stop oral hypoglycemic containing Metformin
 - Examples: Glucophage, Riomet, Glumetza, Glucophage XR, Fortamet, Avandamet, Actoplus, Glucovance, Actoplus, Glyburide, Glipizide, Invokamet, Jentadueto, Janumet, Kombiglyze XR, Metaglip, PrandiMet, Xigduo, Kazano, Synjardy, Segluromet
 - ➤ All other **oral hypoglycemic agents** should **be held** the morning of surgery
 - ➤ Wearable continuous blood glucose monitoring systems: please notify your surgeon and/ or anesthesiologist as soon as possible. GBMC nursing staff will monitor your blood glucose per hospital policy.
 - ➤ If you have a wearable insulin pump system, please notify your surgeon and/or anesthesiologist as soon as possible.
 - ➤ If you are Diabetic notify your endocrinologist of your upcoming surgery
 - ➤Insulin: your Endocrinologist will advise you on how many units they want you to take and when to take give your last dose before surgery.



Diabetes Medications...continued

GLP-1 Agonist:

- semaglutide (*Ozempic*, *Rybelsus*), dulaglutide (*Trulicity*), exenatide (*Bydureon BCise*, *Byetta*), liraglutide (*Victoza*, *Saxenda*), lixisenatide (*Adlyxin*), etc.
- **off label use for weight loss
- ▶ If taken daily: Hold GLP-1 agonists on the day of the procedure/surgery.
- ➤ If taken weekly: Hold GLP-1 agonists a week prior to the procedure/surgery.
- A <u>CLEAR LIQUID ONLY DIET</u> starting *12 hours before the usual NPO time* (noon the day before surgery if NPO).

Clear Liquids for 24h before surgery

- Coffee and tea without milk or non-dairy creamer (sweetener is ok)
- Clear, nonfat broths
- Fruit & vegetable juices that are strained & pulp free
- Sodas & Sports drinks
- Pulp-free popsicles
- Jell-O
- Nothing to eat/drink after 12-midnight



The Night Before Surgery

- ➤ NO heavy meals past 8 pm
 - You can snack up until midnight
- ➤ NO eating past midnight
 - ➤ Including any mints, hard candies, and gum
- > Patients who have the following must STOP liquids 6 hours before surgery:
 - Diabetes
 - Gastroesophageal Reflux Disease (GERD)
 - ➤ BMI >39 (morbidly obese)
 - Gastroparesis (slow moving gut)
- ➤ If you do not have diabetes, GERD, obesity, or gastroparesis:
 - ➤ You CAN drink clear liquids (a max total of 16 ounces) up to 2 hours before your surgery



The Night Before Surgery

- > NO heavy meals past 8 pm
 - You can snack up until midnight
- NO eating past midnight
 - Including any mints, hard candies, and gum
- Patients who have the following must STOP liquids 6 hours before surgery:
 - Diabetes (Type I or Type II)
 - Gastroesophageal Reflux Disease (GERD)
 - > BMI >39 (clinical obese)
 - Gastroparesis (slow moving gut)
- ➤ If you DO NOT have diabetes, or severe acid reflux, or are considered clinically obese (BMI >39), and/or have gastroparesis...then you are safe CAN drink clear liquids (a max total of 16 ounces) up to 2 hours before your surgery time.





Current Medications: Morning of Surgery

Blood pressure/cardiac medication

- ➤ Take your Beta blocker and/or Calcium Chanel blocker heart medication and talk to your primary care or cardiologist to discuss which medications you should take the with a sip of water the morning of your surgery.
- ➤ Discuss with your primary care or cardiologist if they want you to take your diuretics the morning of your surgery
 - ➤ Examples: HCTZ, Lasix

Heartburn or ulcer medicine

- ➤ Acid blockers should be taken on the morning of the surgery to reduce the risk of aspiration pneumonia.
 - Examples: Zantac, Pepcid, Axid, Prilosec, Pantoprazole, Reglan
- ➤ Antacids should NOT be taken they contain particulate material that may damage the lungs if aspirated
 - ➤ Examples: Maalox, Tums, Carafate

No CBD/THC products on the day of your surgery.





What to Bring to the Hospital

- ☐ Your ID card
- Insurance card
- Advance directive (a copy if you have one)
- Preop CHG wash form
- Your cane (if you have one)
- ☐ Bring in glasses, contact lenses, and hearing aids
- □ Regular clothing
 - ➤ Loose fitting clothes that are easy to put on: a tank top, a loose-fitting shirt, or one that buttons up in the front, elastic waist pants or shorts, under garments
 - Shoes you can easily slip onto your feet
- ☐ Any braces, shoe inserts or splints that you normally use
- CPAP machine** (Patients with sleep apnea, overnight stay)





What to Bring to the Hospital

☐ Your ID card ☐ Insurance card(s) ☐ Advance directive (if you have one, bring a copy) ☐ Immobilizer Sling ☐ Cane or other DME equipment from home (place your full name on the device) ☐ Bring your eyeglasses, contact lenses, and hearing aids ☐ Regular clothing ➤ Shirt should be loose fitting through the sleeves and shoulders and button up the front, under garments Walking shoes with good ankle support (no open heel shoes) that is easy to slip into. Do not buy new shoes: bring something comfortable ☐ Any braces, shoe inserts or splints that you normally use

□ CPAP machine** (Patients with sleep apnea, *overnight stay only*)





Visitor Policy: Inpatient Adult Units

- ▶4 visitors are allowed at a time, 1-visitor can stay overnight (9p-8a).
- ➤ Visitors 10-17 years must be accompanied by an adult over 18 yrs. Visitors under 10yrs are not allowed.



- ➤ Visiting Hours 8a-9p
- ➤ Visitors must not be within the first 10 days of positive Covid-19 test
- ➤ Visitors with any respiratory symptoms must wear a mask covering the nose and mouth at all times
- ➤ Visitors must remain at the bedside and use the call-bell for all patient requests
- ➤ If you require an ADA exception, please call the Joint & Spine Center



CENTER



Arriving to the Hospital on the Day of Surgery




- Arrive at the hospital at least 2 hours before your surgery time.
- > Park in Lily Garage.
- Take elevator to the main floor to enter Lobby E.
- ➤ Lobby E is on the 3rd floor of main hospital.





Directions to the General Operating Room (GOR)

➤ Once inside, **Lobby E**, walk down the corridor, you will pass a sitting area on the LEFT.



- Continue until you reach elevator E.
- ➤ Take **Elevator E** on the left to the 4th floor



When you exit the elevator walk straight forward to the GOR registration desk.







Arriving to the Hospital

- Arrive at the hospital at least 2 hours before your surgery time
- ➤ Park in Lily Garage. Enter Lobby E on the 3rd floor of main hospital. Take Elevator E to 4th floor, check in at admission desk









The Pre-operative Area

In Pre-Op

- You will be given a hospital gown to change into
- You will be given in this pre-op area:
 - CHG wipes to clean your skin a 5th time
 - Nozin Nasal Sanitizer to begin application
- Meet your surgical team: surgeon, nurse, anesthesiologist, tech
- Admission database will be completed
- Vital signs, IV will be started; IV fluids and antibiotic(s)







Pre-op Infection Prevention



Do you have an orange fruit oil allergy?

**This product will be given to you the morning of surgery in GBMC's Pre-op Area.







After Surgery


PACU I (postanesthesia care unit)

- > You will wake up and probably feel sleepy
- ➤ Your vital signs will be assessed every 15 minutes
- ➤ Cardiac and oxygen monitors will be in place
- ➤ Pain should be well managed with the nerve block and cocktail, but additional pain relievers are available as needed
- ➤ Your anesthetist will evaluate your progress, once you have met the criteria for the next phase: vital signs are stable, pain is managed, you have sensation in operative leg, and the room you will be transferred to is clean:
 - GBMC's 23-hour/admit patient will be transferred to Unit 58



Hospital Admission: 23-hour observation/ in-patient admit

- Once your vital signs are stable, pain is managed, and you have sensation in operative leg(s). Your anesthetist will clear you for transfer from the PACU to Unit 58.
- Unit 58 is located on the 5th floor of
 Zone D (Labor & Delivery entrance)





Admission to Unit 58

- Private room
- Meet your RN (Registered Nurse) and NST (Nurse Support Tech)
- Communication Board updated
- Ice wrap, Welcome Folder, and your Nozin 12 mL bottle kit will be given to you when you arrive to U58.
 - Upon arrival, the nurse will be teaching you how to give yourself a dose of Nozin from your home kit.





Post-op Infection Prevention

- First dose: after surgery upon arrival to U58
- Next dose: at 9 pm, then every 12 hours
- Continue every 12 hours until your 1st post-op visit or until the bottle is empty (30 days)





Nozin® Nasal Sanitizer® Multidose 12ML Bottle How to Apply

Clean nostrils with a tissue. Discard.



1. Shake



Shake bottle well. Remove cap.


2. Saturate

Apply four (4) drops of solution onto cotton swab tip.

3. Apply

Insert swab tip into RIGHT nostril. Swab eight (8) times clockwise and eight (8) times counterclockwise.

4. Swab



Ensure swab rotation covers all surfaces, including the inside tip of the nostril.

5. Saturate Again



Re-saturate the swab tip by applying **two (2) drops** of solution.

counterclockwise. 6. Apply Left Nostril



Repeat application steps 3 - 4 in the LEFT nostril.

Do not use if you have allergy to citrus oil, or you have nasal bleeding or irritation.

Intended Use: Nozin® Nasal Sanitizer® is a topical antiseptic used to decrease bacteria on the nasal vestibule skin.

Active Ingredient: Alcohol 62%

Precautions: Do not extend applicator into nose beyond swab tip.

Please also refer to Instructions for Use (IFU), product package and Nozin.com.

INOZITI.

62017-2020 Global Life Technologies Corp. Ali rights, reserved. Made in USA Nozin*, Nasal Sanities. Popswab* are trademarks of Global Life Technologies Corp. Patent Noz.: no zin. com joatents

Do you have an *orange fruit oil allergy*?



Fall Risk

- Red non-skid socks will be placed on you, signaling that you are at a high risk for experiencing a fall
- Do NOT get up without the help of staff assisting you.
- Use your call bell









Sequential Compression Device

- Used on your lower extremities to help circulation
- > Help prevent blood clots






Pain Management

- ➤ Pain scale 0-10
 - > "0" means no pain
 - ➤ "10" means the worst pain that you have ever had
- Remember: You should not expect your pain level to be "0" after surgery
- Pain medications are available as needed (prn)

Call your nurse when you are experiencing pain above your goal





Ice Wraps after Surgery

- Placed on your operative joint after surgery
- Decreases pain levels and swelling





Rehabilitation: PT versus OT

Physical Therapy: helps you with functional mobility

- Walking
- Getting in/out of bed
- > Transfers into/out of a chair
- Going up/down steps
- Doing your exercise program

Occupational Therapy: helps you with functional activities of daily living (ADL)

- Getting dressed and bathed
- ➤ Getting in/out of tub/shower
- Using the toilet
- Car transfers
- How to use adaptive equipment if needed
- Getting items out of refrigerator/cabinets
- Show you how to take your sling on and off



Physical and Occupation Therapy: Their Assessment and Education

- Social information: Do you live alone or have help?
- Environment during recovery:
 - > Do you live on one level or more?
 - Number of steps inside/outside home?
 - ➤ Are there railings?
 - ➤ Bathtub or shower equipment?
 - > Take pictures of your bathroom, kitchen, living room, etc.
- Prior level of function: How were you doing before you came into the hospital?



Immobilizer Sling

- After surgery, you will be placed in an immobilizer sling to limit the movement of your shoulder.
- You will wear the sling at all times except when showering or doing approved exercises shown to you by the therapy team





Medical Equipment Needs

Your OT will assess, evaluate, and determine what assistive devices may needed at home

- **≻**Cane
- **≻**Reacher
- ➤ Long handled shoehorn
- ➤ Long handled sponge
- > Elastic shoelaces
- ➤ Sock aid













GBMC Care Management



The Care Manager (CM) on Unit 58 will visit you the day after your surgery and follow the inpatient therapist's recommendation for therapy. The CM will continue working on your discharge planning needs, offer you resources, and communicate the plan to your care team.

- ➤ Home with Out-Patient Therapy: Your GBMC occupational therapist (OT) and/or physical therapist (PT) will make recommendations for therapy needs. Most patients begin outpatient therapy two weeks post-operation. You will be given an outpatient prescription at the time of your discharge.
 - ➤ Designate who is driving you to your appointment
- Equipment for home: OT/PT will evaluate if you need any DME, your case manager will contact your insurance company for approval and arrange delivery to your room.
- Skilled Nursing Facility (SNF): Medical necessity and insurance authorization approval needed. If your insurance denies discharge to a subacute or acute rehab center, you will be discharged home.
- ➤ Discharge is between 1pm-5pm

Pain Medication at Discharge

- ➤ PICK-UP your pain medications a week before surgery! Your surgeon's office will send your prescriptions electronically to your preferred pharmacy. Please let your surgeon know if your pharmacy location has changed.
- ➤ Could be prescribed a 7-day supply of medications
 - ➤ Dependent upon your insurance
 - Verify with your pharmacy at time of pick up
- ➤ Per the Maryland Department of Health & Mental Hygiene guidelines, you will be prescribed an opioid antidote called Naloxone (Narcan) in the event of an overdose
- ➤ Call the office three days in advance of the need to have medications renewed
- ➤ No prescriptions will be addressed on weekends or after hours
- ➤ Pain medications and/or muscle relaxers will only be prescribed for up to 3 months following your surgery
 - ➤ After that period of time: if needed, you will be referred to a pain management specialist





Discharge Day

- ➤ GBMC Same-Day Outpatient surgery, patients will be discharged the day of surgery from GBMC's PACU recovery area
- ➤ In-patient surgery are discharged the day after their surgery
 - Most patients are discharged after 12 pm, when your:
 - > PT and OT have both cleared you for discharge
 - A Case Manager has visited you
 - A cane (if needed) is delivered to your room
 - You are given your prescription for medications, if you were not already given them by your surgeon's office before surgery, (can be electronically sent to your pharmacy)
 - Nurse has given you your discharge paperwork
 - Your support person has arrived to the hospital



Discharge Day Checklist Guide- Shoulder I have reviewed my discharge instructions with the nurse ☐ I know who is my support person/help at home ☐ Post-op Physical Therapy Agency (if needed): _____ Phone # > If I haven't heard from the Physical Therapy Agency 3-days after discharge or have any issues with scheduling, I will call my surgeon's office as soon as possible Do I have my prescription(s)? Medication(s) MUST be picked up the day of discharge GBMC Walgreens can deliver your medications to your room, accepts cash or credit cards Do I know the reason for and side effects of my prescriptions? ☐ I have my... Cane (if needed) Sling Ice packs with the wrap Dressing material (gauze and tape) ➤ _____ Nozin® Nasal Sanitizer® 12-mL bottle and starter cotton swabs Belongings that I brought into the hospital I will call my surgeon's office with any signs of infection such as fever, redness, swelling, tenderness, or puss-like drainage I will call my surgeon's office with any new loss or decrease in sensation in the hands or fingers of the operative arm Please contact your surgeon's office with any questions at (

Please review the
Joint & Spine Center
Pre-op Packet
for important information,
including this
post-op checklist
seen here.





Thank you for choosing GBMC









Joint and Spine Center 6701 N Charles St Towson, MD 21204

Phone: 443-849-6261

Email: jointspinecenter@gbmc.org

Website: www.gbmc.org/JointandSpine

ACTIVATE your MyChart today







CENTER